



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : B01J 20/32, C02F 1/28, C07K 1/22, C07D 251/50, 251/54, 403/12, 251/70, A61M 1/36, G01N 33/50	A1	(11) International Publication Number: WO 00/67900 (43) International Publication Date: 16 November 2000 (16.11.00)
(21) International Application Number: PCT/GB00/01759 (22) International Filing Date: 9 May 2000 (09.05.00) (30) Priority Data: 9910807.8 10 May 1999 (10.05.99) GB (71) Applicant (for all designated States except US): PROMETIC BIOSCIENCES LTD. [GB/GB]; Freeport, Ballasalla, Isle of Man IM9 2AP (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): LOWE, Christopher, Robin [GB/GB]; Institute of Biotechnology, University of Cambridge, 20 Tennis Court Road, Cambridge CB2 1QT (GB). LAWDEN, Kim, Hilary [GB/GB]; 8 Johnson Mansions, Queens Club Gardens, West Kensington, London W14 9SJ (GB). (74) Agent: GILL JENNINGS & EVERY; Broadgate House, 7 Eldon Street, London EC2M 7LH (GB).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NOVEL TRIAZINE-BASED DETOXIFICATION AGENTS AND THEIR USE <div style="text-align: center;"> <p style="text-align: right;">(I)</p> </div> (57) Abstract <p>An affinity ligand-matrix conjugate comprises the matrix and, conjugated thereto by the group Z, a ligand having general formula (I) wherein one X is N and the other X is N, CCL or CCN; A₁ and A₂ are each independently O, S or N-R₁ and R₁ is H, C₁₋₆alkyl, C₁₋₆ hydroxyalkyl, benzyl or β-phenylethyl; B₁ and B₂ are each independently an optionally substituted hydrocarbon linkage containing from 1 to 10 carbon atoms; D₁ is H or a primary amino, secondary amino, tertiary amino, quaternary ammonium, imidazole, guanidino or amidino group; and D₂ is a secondary amino, tertiary amino, quaternary ammonium, imadazole, guanidino or amidino group; or B₂-D₂ is -CHCOOH-(CH₂)₃₋₄-NH₂; AND p is 0 or 1. Such conjugates are useful for the separation, isolation, purification, characterisation, identification or quantification of an endotoxin.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NOVEL TRIAZINE-BASED DETOXIFICATION AGENTS AND THEIR USE FIELD OF THE INVENTION

The present invention relates to novel affinity ligands, their preparation and attachment to matrices which may consist of solid, semi-solid, particulate or colloidal materials, or soluble polymers. The invention furthermore relates to these novel affinity ligand-matrix conjugates and the preparation and use thereof in the binding and removal of endotoxin from various fluids such as water, aqueous solutions, body fluids, blood, plasma, solutions of pharmaceutical products, proteins and other compounds of biological origin.

BACKGROUND OF THE INVENTION

Endotoxins are lipopolysaccharides found in the outermost membrane of Gram-negative bacteria, particularly pathogenic bacteria of the class Enterobacteriaceae, Neisseriaceae and Chlamydiaceae. Endotoxins comprise lipid A attached to a polysaccharide of variable structure dependent upon its biological origin. The polysaccharide component of Enterobacteriaceae endotoxin is characterised by an O-specific chain region and a core region. The O-specific region comprises up to 50 repeating oligosaccharide units that contain as many as 8 different sugar residues. O-specific chains exhibit large structural diversity from species to species whereas the core region, divided into the outer core and inner core regions, is less variable. The inner core region is characterised by the presence of unusual sugar residues such as heptose and 2-keto-3-deoxyoctonic acid (KDO) which are frequently substituted with phosphate or phosphate derivatives. Also attached to the inner core region, lipid A is a conserved biphosphorylated glucosamine disaccharide which is acylated by 4 saturated primary acyl groups of which 2 carry secondary saturated acyl groups. The combination of hydrophobic lipid A tails with the hydrophilic and anionic polysaccharide unit provides endotoxin with amphipathic properties.

Endotoxin released from the cell wall of Gram-negative bacteria is considered to be the primary cause of the many pathophysiological occurrences that accompany Gram-negative septicemia. Endotoxin at pg/ml concentrations in blood triggers the release of a variety of cytokines, including interleukins and TNF. Over stimulation of the immune system by endotoxin leads to a massive

release of cytokines which ultimately results in metabolic breakdown and septic shock. During septic shock, the complement and coagulation cascades become activated and vascular permeability increases. This can lead to disseminated intravascular coagulation and multiple organ failure, often with fatal consequences. Septic shock often develops because of the lack of an initial response to infection allowing the level of blood-borne endotoxin to reach critical levels.

In addition to the obvious risk presented by the presence of live Gram negative bacteria or cell wall debris in parenteral pharmaceutical products, the presence of free endotoxin in pharmaceutical preparations is also a major concern. Because endotoxin is such a potent immune stimulator, very low concentrations may cause toxic reactions including pyrogenic effects. Endotoxin is a relatively stable molecule which is not inactivated by routine autoclaving or treatment with organic solvents. Exposure to concentrated sodium hydroxide or prolonged high temperature (250°C) will inactivate endotoxin, though such methods are not appropriate for most biological products. Furthermore, maintenance of complete sterility throughout the manufacture of bio-therapeutics is problematic. Consequently, the highly efficient capture and removal of endotoxin from parenteral pharmaceuticals is very desirable, particularly in situations where endotoxin is known to associate with components of the therapeutic formulation.

A variety of techniques have been used to remove endotoxin from aqueous solutions including ultrafiltration, charcoal adsorption, cation-exchange chromatography, and a variety of immobilised affinity ligands including polymyxin B and endotoxin binding protein. All of these techniques exhibit significant shortcomings, particularly in the case of endotoxin removal from high molecular weight compounds such as therapeutic proteins. Ultrafiltration can only be used to remove endotoxin from low molecular weight compounds whereas charcoal adsorption tends to promote the binding of most organic compounds. Cation-exchange chromatography is effective in removing endotoxin from water but less effective for protein containing solutions, particularly proteins with acidic isoelectric points. Polymyxin B, a cyclic polypeptide antibiotic, is too toxic to

allow its use for the purification of therapeutic products whereas endotoxin binding protein is too expensive for commercial applications.

Immobilised cationic amino acids (histidine, lysine and arginine) have also been used for endotoxin removal (Tosa, T. *et al.*, Molecular Interactions in Bioseparations, Ed. Ngo, T. T., Plenum Press, New York, pp. 323-332, 1993; Lawden, K. H. *et al.*, Bacterial Endotoxins: Lipopolysaccharides From Genes to Therapy, Wiley-Liss Inc., pp. 443-452, 1995). Such materials have been prepared by direct attachment of amino acids to epoxy-activated chromatographic matrices. In the case of Pyrosep™, a commercially available material manufactured by Tanabe Seiyaku Company Limited, Osaka, Japan, a single histidine group is immobilised to a support matrix by a hexanediamine spacer arm. Again, such materials are adequate for removal of endotoxin from water or solutions of low molecular weight compounds, but their performance is compromised in the presence of salt (>50 mM) or proteins which have an affinity for endotoxin. Consequently, none of the existing methods of endotoxin removal are suited to the elimination of endotoxin from bio-therapeutic compounds intended for parenteral administration. This is especially true for protein therapeutics where no single effective and safe method of endotoxin removal exists.

Removal of endotoxin from blood or plasma may provide an effective approach to the management of septic shock, particularly if applied at the early stages of infection or prophylactically in situations where an increased risk of septic shock is anticipated (e.g. major bowel or liver surgery). Several studies have been reported as to the use of monoclonal antibodies directed against endotoxin or cytokines released in the initial phase of the shock reaction. However, most of these approaches have been found to be ineffective (Siegel, J. P., Drug Information Journal, 30, pp. 567-572, 1996). In contrast, extracorporeal extraction of endotoxin from whole blood has been accomplished by use of fibre-immobilised polymyxin B (Aoki, H. *et al.*, Nippon Geka Gakkai Zasshi (Japan), 94, pp. 775-780, 1993), though concerns over potential toxicity of polymyxin B leachates remain. Consequently, affinity adsorbents incorporating

endotoxin binding ligands which have high affinity for endotoxin and low toxicity may also be beneficial for the management of sepsis.

Immobilised amino acids have also been investigated as potential endotoxin removal agents but such materials bind endotoxin weakly and non-specifically and are of limited value in the extraction of endotoxin from biological fluids and solutions of biological compounds. Triazine-based compounds have been reported which bind selectively to proteins; however, such ligands are not applicable to the isolation of endotoxin.

SUMMARY OF THE INVENTION


This invention relates to the discovery of synthetic affinity ligand structures which bind selectively to endotoxin. A generic group of novel affinity ligands have been found which exhibit high affinity for endotoxin and are generally applicable to the isolation of endotoxin from a variety of sources.

A feature of the present invention is the provision of a general tool for the removal of endotoxin contamination from biological materials. Endotoxin binds exceedingly tightly to affinity ligand-matrix conjugates of the invention. This feature enables highly efficient extraction of endotoxin from water and aqueous solutions providing a means of generating pyrogen-free water or pyrogen-free solutions. Affinity ligand-matrix conjugates of the invention are especially valuable for the removal of endotoxin which is bound to or associated with proteins, drugs or other biological compounds intended for medical or pharmaceutical applications. Certain biological compounds, particularly proteins, often bind endotoxin tightly and subsequent removal is very difficult, if not impossible, by existing means. Affinity ligand-matrix conjugates of the invention may also be applied to the removal of endotoxin from blood or plasma and so provide an especially useful tool for *in vitro* or *in vivo* removal of endotoxin, the latter being achieved, for example, by way of an extracorporeal endotoxin extraction device. Such a device may be especially valuable for removal of endotoxin which is released into the blood stream during bacterial infections, such infections often causing life-threatening diseases such as septicaemia or meningitis. Removal of blood-borne endotoxin may be particularly beneficial in

the treatment of these diseases and in the prevention and management of septic shock.

Novel affinity ligand-matrix conjugates provided by this invention can be used in place of other endotoxin binding materials and are significantly more flexible in their use, are more robust, less expensive to produce and offer greater endotoxin binding efficiencies.

The present invention relates to affinity ligand-matrix conjugates comprising a ligand having General Formula (1):



10 wherein one of the symbols X represents a nitrogen atom and the other symbol X represents a nitrogen atom or a carbon atom carrying a chlorine atom or a cyano group;

A₁ and A₂ each independently represent an oxygen atom, a sulphur atom or a group N-R_i;

15 R₁ represents a hydrogen atom, an alkyl group containing from 1 to 6 carbon atoms, a hydroxyalkyl group containing from 1 to 6 carbon atoms, a benzyl group or a β-phenylethyl group;

B₁ and B₂ each independently represent an optionally substituted hydrocarbon linkage containing from 1 to 10 carbon atoms (any substituent being substantially non-critical with respect to utility) and including alkyl, phenyl, naphthyl and cyclohexyl groups;

D₁ represents a hydrogen atom, a primary amino group, a secondary amino group, a tertiary amino group, a quaternary ammonium group, an imidazole group, a guanidino group or an amidino group;

D₂ represents a primary amino group (e.g. as derived from lysine or ornithine), a secondary amino group, a tertiary amino group, a quaternary ammonium group, an imidazole group, a guanidino group or an amidino group; and

p is 0 or 1.

The ligand is attached to a support matrix in position Z, optionally through a spacer arm interposed between the ligand and matrix. Alternatively, in novel ligands of the invention, Z represents a functional group of the type capable of reaction with a solid matrix that may be activated (if necessary or desired) or unactivated.

DESCRIPTION OF THE INVENTION

When conjugated to a matrix, the optional spacer arm is preferably represented by General Formula (II):



wherein T represents an oxygen atom, a sulphur atom or a group N-R₂; wherein R₂ represents a hydrogen atom or an alkyl group containing from 1 to 6 carbon atoms;

V represents an oxygen atom, a sulphur atom, a -COO- group, a CONH group or an NHCO group, a -PO₃H group, a NH-arylene-SO₂-CH₂-CH₂-group or a N-R₃ group; wherein R₃ represents a hydrogen atom or an alkyl group containing from 1 to 6 carbon atoms;

L represents an optionally substituted hydrocarbon linkage containing from 2 to 20 carbon atoms; and


m is 0 or 1.

The support matrix may be any compound or material, particulate or non-particulate, soluble or insoluble, porous or non-porous which may be used in conjunction with affinity ligands to form an affinity ligand-matrix conjugate and

which provides a convenient means of separating the affinity ligands from solutes in a contacting solution.

The present invention provides novel affinity ligand-matrix conjugates, which affinity ligand-matrix conjugates may be used in the isolation or removal of endotoxin from water, aqueous solutions, body fluids, blood, plasma, solutions of pharmaceutical products, proteins and other compounds of biological origin.

In a preferred embodiment, the invention provides novel affinity ligand-matrix conjugates which are represented by the General Formula (III):



wherein A_1 , A_2 , B_1 , B_2 , D_1 , D_2 , p , X , T , L , V , m , R_1 , R_2 , and R_3 have the meanings specified above; and M represents the residue of a support matrix which may be any compound or material, particulate or non-particulate, soluble or insoluble, porous or non-porous which may be used in conjunction with affinity ligands to form an affinity ligand-matrix conjugate and which provides a convenient means of separating the affinity ligands from solutes in a contacting solution.

It will be appreciated that this invention relates, inter alia, to the use of compounds which are pyrimidines, diazines, or triazines carrying a $-T-[L-V]_m-M$ substituent, or the precursor thereof, and other substituents linked to the ring via a hetero atom. Such substituents may include any non-interfering group comprising 0 to 20 carbon atoms.

In the present specification, whenever the term endotoxin is used in a plural or generic sense, it is intended to mean endotoxins originating from any microbiological source. By the term endotoxin is thus also meant lipopolysaccharide from any species including Enterobacteriaceae, 5 Neisseriaceae and Chlamydiaceae. Since endotoxin is known to be heterogeneous, the term "endotoxin" as used herein includes all naturally occurring forms which comprise lipid A covalently linked to a polysaccharide, including analogues, derivatives, fragments and precursors thereof.

The term "primary amino group" as used herein, alone or in combination, 10 refers to an $-NH_2$ group.

The term "secondary amino group" as used herein, alone or in combination, refers to a $-NHR_4$ group; wherein R_4 represents a straight or branched alkyl group containing from 1 to 6 carbon atoms.

The term "tertiary amino group" as used herein, alone or in combination, 15 refers to a $-NR_5R_6$ group; wherein R_5 and R_6 each represent a straight or branched alkyl group containing from 1 to 6 carbon atoms.

The term "quaternary ammonium group" as used herein, alone or in combination, refers to a $-NR_7R_8R_9^+$ group; wherein R_7 , R_8 and R_9 each represent a straight or branched alkyl group containing from 1 to 6 carbon atoms.

20 The term "alkyl group containing from 1 to 6 carbon atoms" as used herein, alone or in combination, refers to a straight or branched, saturated hydrocarbon chain having 1 to 6 carbon atoms such as e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, 2-methylbutyl, 3-methylbutyl, n-hexyl, 4-methylpentyl, neopentyl and 2,2-dimethylpropyl.

25 The term "hydroxyalkyl group containing from 1 to 6 carbon atoms" as used herein, alone or in combination, refers to a straight or branched, saturated hydrocarbon chain having 1 to 6 carbon atoms substituted with one or more hydroxy groups, preferably one hydroxy group, such as e.g. hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, 4-hydroxybutyl, 5-hydroxypentyl 30 and 6-hydroxyhexyl.

The term "alkoxy group containing from 1 to 6 carbon atoms" as used herein, alone or in combination, refers to a straight or branched monovalent

substituent comprising an alkyl group containing from 1 to 6 carbon atoms linked through an ether oxygen having its free valence bond from the ether oxygen and having 1 to 6 carbon atoms e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy and pentoxy.

5 The term "halogen" means fluorine, chlorine, bromine or iodine.

 The term "acyloxy or acylamino containing from 1 to 6 carbon atoms" as used herein refers to a monovalent substituent comprising an alkyl group containing from 1 to 5 carbon atoms linked through a carbonyloxy or oxycarbonyl group such as a methylcarbonyloxy, ethylcarbonyloxy, methyloxycarbonyl or
10 ethyloxycarbonyl group or linked through a carbonylamino or aminocarbonyl group such as a methylcarbonylamino, ethylcarbonylamino, methylaminocarbonyl or ethylaminocarbonyl group.

 The term "alkysulfonyl containing from 1 to 6 carbon atoms" as used herein refers to a monovalent substituent comprising an alkyl group containing
15 from 1 to 6 carbon atoms linked through a sulfonyl group such as e.g. methylsulfonyl, ethylsulfonyl, n-propylsulfonyl, isopropylsulfonyl, n-butylsulfonyl, sec-butylsulfonyl, isobutylsulfonyl, tert-butylsulfonyl, n-pentylsulfonyl, 2-methylbutylsulfonyl, 3-methylbutylsulfonyl, n-hexylsulfonyl, 4-methylpentylsulfonyl, neopentylsulfonyl, and 2,2-dimethylpropylsulfonyl.

20 The term "one or more substituents independently selected from" shall more preferably refer to from 1-3 substituents. The term shall further preferably refer to 1-2 substituents and most preferably refer to one substituent.

 The term "optionally substituted hydrocarbon linkage containing from 2 to 20 carbon atoms" as used herein refers to one or more linear or branched alkyl
25 chains, optionally substituted with for example hydroxy or alkoxy groups containing from 1 to 6 carbon atoms, and optionally linked together by amino, ether, thioether, ester, amide or sulphonamide bonds providing a chain containing from 2 to 20 carbon atoms. The construction is preferably flexible. The construction of such optionally substituted hydrocarbon linkages is for
30 example described in Lowe, C.R. and Dean, P.D.G, 1974, Affinity Chromatography, John Wiley & Sons, London, which is hereby incorporated by reference.

The term "optionally substituted hydrocarbon linkage containing from 1 to 10 carbon atoms" as used herein, alone or in combination, refers to a linear or branched hydrocarbon chain having 1 to 10 carbon atoms optionally substituted with one or more functional groups, including but not limited to, carboxyl groups, preferably one carboxyl group, and hydroxyl groups.

In a preferred embodiment of the invention, R_1 represents a hydrogen atom.

In another preferred embodiment of the invention, R_2 represents a hydrogen atom.

In another preferred embodiment of the invention, R_3 represents a hydrogen atom.

In another preferred embodiment of the invention, R_4 represents a methyl group, an ethyl group or a propyl group.

In another preferred embodiment of the invention, R_5 represents a methyl group, an ethyl group or a propyl group.

In another preferred embodiment of the invention, R_6 represents a methyl group, an ethyl group or a propyl group.

In another preferred embodiment of the invention, R_7 represents a methyl group, an ethyl group or a propyl group.

In another preferred embodiment of the invention, R_8 represents a methyl group, an ethyl group or a propyl group.

In another preferred embodiment of the invention, R_9 represents a methyl group, an ethyl group or a propyl group.

In another preferred embodiment of the invention, A_1 represents $N-R_1$ wherein R_1 is as defined above.

In another preferred embodiment of the invention, A_2 represents $N-R_1$ wherein R_1 is as defined above.

In another preferred embodiment of the invention, B_1 represents a -CHCOOH-CH₂-group, a -CHCOOH-(CH₂)₂-group, a -CHCOOH-(CH₂)₃-group, a -CHCOOH-(CH₂)₄-group, an ethyl group, a propyl group, a 2-hydroxypropyl group, a butyl group, a pentyl group, a hexyl group or a phenyl group.

In another preferred embodiment of the invention, B_2 represents a

-CHCOOH-CH₂- group, -CHCOOH-(CH₂)₂- group, a -CHCOOH-(CH₂)₃- group, a -CHCOOH-(CH₂)₄- group, ethyl group, a propyl group, a 2-hydroxypropyl group, a butyl group, a pentyl group, a hexyl group or a phenyl group.

In another preferred embodiment of the invention, D₁ represents
5 hydrogen, an amino group, an imidazole group, a guanidino group, an aminidino group, a trimethylammonium group, a triethylammonium group, a dimethylamino group, a diethylamino group, a methylamino group or an ethylamino group.

In another preferred embodiment of the invention, D₂ represents an amino group, an imidazole group, a guanidino group, an aminidino group, a
10 trimethylammonium group, a triethylammonium group, a dimethylamino group, a diethylamino group, a methylamino group or an ethylamino group. D₂ (and often also D₁) is preferably a strongly charged species.

In another preferred embodiment of the invention, p represents 0 or 1.

In another preferred embodiment of the invention, both X represent a
15 nitrogen atom.

In another preferred embodiment of the invention, T represents an oxygen atom or, more preferably, an NH group.

In another preferred embodiment of the invention, L represents a butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl or dodecyl group and V and m are as
20 defined above.


In another preferred embodiment of the invention, V represents an oxygen atom, a -COO- group, a PO₃H- group or an N-R₃ group; and more preferred an oxygen atom or an NH group and L and m are as defined above.

In another preferred embodiment of the invention, m represents 1 and L
25 and V are as defined above.

The term "integer between x and y" may include the values x (including zero) and y.

The invention also provides methods for the manufacture of novel affinity ligand-matrix conjugates according to the invention which comprises reacting,
30 in any order,

(i) a halogenoheterocyclic compound of General Formula (IV):



wherein the symbols X have the meaning hereinbefore specified and W represents a halogen atom with

5 (ii) a compound of General Formula (V):



wherein the symbols D_1 , B_1 , A_1 and p have the meanings hereinbefore
10 specified and H is hydrogen,

(iii) a compound of General Formula (VI):



15

wherein the symbols D_2 , B_2 and A_2 have the meanings hereinbefore specified and H is hydrogen, and

(iv) with either an optionally derivatised support matrix of General Formula
20 (VII):



wherein the symbols T, L, V, m and M have the meanings hereinbefore specified and H is hydrogen


or, with a linking unit of General Formula (VIII):

5



wherein the symbols T, L, V have the meanings hereinbefore specified to give a compound of General Formula (IX):

10



wherein A₁, A₂, B₁, B₂, D₁, D₂, p, X, T, L, V, m, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ have the meanings hereinbefore specified; the compound of General Formula (IX) is then reacted further with a support matrix whose residue is represented by M using activating and coupling procedures well known to those skilled in the art.

As examples of halogenoheterocyclic compounds of General Formula (IV) there may be mentioned 5-chloro-2,4,6-trifluoropyrimidine, 5-cyano-2,4,6-trichloropyrimidine, cyanuric fluoride, cyanuric bromide and, above all, cyanuric chloride.

As examples of compounds of General Formula (V) there may be mentioned ammonia, water, arginine, lysine, histidine, α,γ-diaminobutyric acid, m-aminobenzamidine, p-aminobenzamidine, m-aminobenzenetrimethyl-

ammonium bromide, p-aminobenzenetrimethylammonium bromide, 2-(diethylamino)ethylamine, (2-aminoethyl)trimethylammonium chloride, histamine, agmatine, ethylenediamine, 1,3-diaminopropane, 1,3-diamino-2-hydroxypropane, 1,4-diaminobutane, 1,5-diaminopentane and 1,6-diaminohexane.

5 As examples of compounds of General Formula (VI) there may be mentioned arginine, lysine, histidine, α,γ -diaminobutyric acid, m-aminobenzamidine, p-aminobenzamidine, m-aminobenzenetrimethylammonium bromide, p-aminobenzenetrimethylammonium bromide, 2-(diethylamino)ethylamine, (2-aminoethyl)trimethylammonium chloride, histamine,
10 agmatine, ethylenediamine, 1,3-diaminopropane, 1,3-diamino-2-hydroxypropane, 1,4-diaminobutane, 1,5-diaminopentane, 1,6-diaminohexane.

As example of support matrices whose residue is represented by M, there may be mentioned insoluble support matrices such as a naturally occurring polymer, for example a polypeptide or protein such as cross-linked albumin or
15 a polysaccharide such as agarose, alginate, carrageenan, chitin, cellulose, dextran or starch; synthetic polymers such as polyacrylamide, polystyrene, polyacrolein, polyvinyl alcohol, polymethylacrylate or perfluorocarbon; inorganic compounds such as silica, glass, kieselguhr, alumina, iron oxide or other metal oxides or co-polymers consisting of any combination of two or more of a naturally
20 occurring polymer, synthetic polymer or inorganic compounds. Also included within the definition of support matrices whose residue is represented by M are soluble support matrices comprising polymers such as dextran, polyethylene glycol, polyvinyl alcohol or hydrolysed starch, which provide affinity-ligand matrix conjugates for use in liquid partitioning; or support matrices comprising
25 compounds such as perfluorodecalin which provide affinity-ligand matrix conjugates for use in the formation of affinity emulsions. For the avoidance of doubt, a support matrix is defined herein as any compound or material whether particulate or non-particulate, soluble or insoluble, porous or non-porous which may be used to form a novel affinity ligand-matrix conjugate according to the
30 invention and which provides a convenient means of separating the affinity ligand from solutes in a contacting solution.

Also included within the definition of support matrices whose residue is represented by M are support matrices such as agarose, cellulose, dextran, starch, alginate, carrageenan, synthetic polymers, silica, glass and metal oxides which have been, or are, modified by treatment with an activating agent prior to, or during, attachment of the ligand.

In a preferred embodiment of the invention, M represents optionally activated agarose, silica, cellulose, dextran, glass, toyopearl, hydroxyethylmethacrylate, polyacrylamide, styrenedivinylbenzene, Hyper D, perfluorocarbons, polysulphone, polyethersulphone, polyvinylidene fluoride, nylon, and polyvinylchloride. Preferably M represents optionally tresyl-activated, sulphonyl chloride-activated, tosyl-activated, vinylsulphone-activated or epoxy-activated agarose.

There exists a considerable number of activating agents which have found use for the general purpose of attaching ligands to support matrices. These compounds and their method of use are well known to those skilled in the art and, since the nub of the present invention lies in the nature of the ligand attached to the matrix and not in the mode of attachment, any of these activating agents will serve in the preparation of the new matrix-ligand conjugates of the invention. As non-limiting examples of such activating agents there may be mentioned such diverse compounds as cyanogen bromide, cyanuric chloride, epichlorohydrin, divinyl sulphone, p-toluenesulphonyl chloride, 1,1'-carbonyldiimidazole, sodium meta-periodate, 2-fluoro-1-methylpyridiniumtoluene-4-sulphonate, glycidoxypolytrimethoxysilane and 2,2,2-trifluoroethanesulphonyl chloride. As indicated above, the procedures by which such activating steps are carried out are well known to those skilled in the art.

Similarly, a wide variety of condensing agents may be used to attach the compounds of General Formulae (VIII) and (IX) to support matrices such as agarose, cellulose, dextran, starch, alginate, carrageenan, silica or glass. Again these compounds, and their method of use are well known to those skilled in the art and, again, since the nub of the present invention lies in the nature of the ligand and not in the mode of attachment, any of these condensing agents will serve in the preparation of the new matrix-ligand conjugates of the invention. As


non-limiting examples of such condensing agents, there may be mentioned N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline, dicyclohexyl carbodiimide and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide.

As examples of linking units of General Formula (VIII) which may be used
5 to produce compounds of General Formula (IX) there may be mentioned diamines such as ethylenediamine, N,N'-dimethylethylenediamine, N-ethylethylenediamine, N-(β -hydroxyethyl)ethylenediamine, propylenediamine, N-methylpropylenediamine, N-(β -hydroxyethyl)propylenediamine, 1,4-diaminobutane, 1,5-diaminopentane, 1,6-diaminohexane, 1,7-diaminoheptane,
10 1,8-diaminooctane, 1,9-diaminononane, 1,10-diaminodecane, 1,12-diamindodecane, piperazine, 3-hydroxy-1,5-diaminopentane, m- and p-phenylene diamine, m- and p-aminobenzylamine; amino alcohols such as ethanolamine, N-methylethanolamine, N-propylethanolamine, diethanolamine, 3-hydroxypropylamine, 2,3-dihydroxypropylamine, isopropanolamine, 5-aminopentan-1-ol and 6-aminohexan-1-ol; aminophenols such as o-, m- and p-aminophenol, aminocarboxylic acids such as glycine, N-methylglycine, 3- and 4-aminobutyric acid, 3-aminoisobutyric acid, 5-aminovaleric acid, 6-aminocaproic acid, 7-aminoheptanoic acid, m- and p-aminobenzoic acid; aminophosphonic acids such as m-aminobenzenephosphonic acid and p-aminobenzylphosphonic
15 acid; and aminoarylene vinylsulphone precursors such as aniline-3- β -sulphatoethylsulphone and aniline-4- β -sulphatoethylsulphone.

The reaction of halogenoheterocyclic compounds of General Formula (IV) with compounds of General Formulae (V), (VI) and (VII) or (VIII) may be carried out in an organic solvent which is not miscible with water; or in an organic
25 solvent which is miscible with water, or in a mixture of water and a water miscible organic solvent. Examples of suitable organic solvents which are not miscible with water are toluene, xylene or chlorobenzene; examples of suitable organic solvents which are miscible with water are acetone, methyl ethyl ketone or dioxan. The first reaction of the halogenoheterocyclic compound may be carried
30 out at temperatures between 0°C and 25°C, ideally between 0°C and 5°C; the second reaction may be carried out at temperatures between 20°C and 50°C, ideally between 30°C and 45°C and the third reaction at temperatures between


20°C and 100°C. During such reactions, the inorganic acid such as hydrochloric acid or hydrofluoric acid which is produced is neutralised by the use of an acid binding agent such as sodium hydroxide, sodium carbonate, sodium bicarbonate, calcium hydroxide or calcium carbonate.

5 Additionally, compounds of General Formula (IX) may be reacted with a reactive polymerisable monomer to form a polymerisable compound of General Formula (X):



wherein A₁, A₂, B₁, B₂, D₁, D₂, p, X, T, L, V, m, R₁, R₂, R₃, R₄, R₅, R₆, R₇, R₈, and R₉ have the meanings hereinbefore specified; R₁₁ represents a hydrogen atom or an alkyl group containing from 1 to 6 carbon atoms; R₁₀ represents a carbonyl group, a methylene group, a -NH-CH₂- group or a -S-CH₂- group. Examples of reactive polymerisable monomers include acryloyl chloride, methacryloyl chloride, allyl bromide, allylamine or 3,4-epoxybutene. Polymerisable compounds of General Formula (X) may be polymerised, optionally in the presence of other polymerisable monomers, to form affinity ligand matrix conjugates of General Formula (III). Such polymerisation procedures are well known to those skilled in the art.


In another embodiment the invention relates to novel affinity ligand matrix
20 conjugates of General Formula (XI):



wherein A_1 , A_2 , B_1 , B_2 , D_1 , D_2 , p , X , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , and R_9 have the meanings hereinbefore specified and Halogen represents a fluorine, chlorine, bromine or iodine atom.

5 Furthermore, the invention relates to a method of attaching novel affinity ligands of General Formula (XI) as defined above to a matrix of General Formula (VII) as defined above by reacting the novel affinity ligands with the matrix at temperatures between 0°C and 100°C, optionally in the presence of an acid binding agent.


10 In another embodiment the invention relates to novel affinity ligands of General Formula (XII):



15 wherein A_1 , A_2 , B_1 , B_2 , D_1 , D_2 , p , X , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , and R_9 have the meanings hereinbefore specified and j is an integer between 2 and 20.

Furthermore, the invention relates to a method of preparing above novel affinity ligands by reacting a compound of above General Formula (XI) with an alkylene diamine of General Formula $H_2N-(CH_2)_j-NH_2$ at temperatures between $0^\circ C$ and $100^\circ C$, optionally in the presence of an acid binding agent.

- 5 In another embodiment the invention relates to novel affinity ligands of General Formula (XIII):



wherein A_1 , A_2 , B_1 , B_2 , D_1 , D_2 , p , X , R_1 , R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , R_8 , and R_9 have the meanings hereinbefore specified; j is an integer between 2 and 20, and q is 0 or

- 10 1.


Furthermore, the invention relates to a method of attaching novel affinity ligands of General Formula (XIII) as defined above to a matrix of General Formula (VII) as defined above by reacting the novel affinity ligands with the matrix at temperatures between $0^\circ C$ and $100^\circ C$ in the presence of a condensing agent. As non-limiting examples of such condensing agents, there may be mentioned N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline, dicyclohexyl carbodiimide and 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide.

- 15 Furthermore, the invention relates to a method of preparing novel affinity ligands of above General Formula (XIII) by reacting a compound of above General Formula $H_2N-(CH_2)_j-(CO)_q-OH$ at temperatures between $0^\circ C$ and $100^\circ C$, optionally in the presence of an acid binding agent.

In another embodiment, the invention relates to novel affinity ligands of above General Formula (X) wherein A_1 , A_2 , B_1 , B_2 , D_1 , D_2 , p , X , T , L , V , m , R_1 ,


$R_2, R_3, R_4, R_5, R_6, R_7, R_8$, and R_9 have the meanings hereinbefore specified; R_{10} represents a hydrogen atom or an alkyl group containing from 1 to 6 carbon atoms; R_{11} represents a carbonyl group, a methylene group, a $-NH-CH_2-$ group or a $-S-CH_2-$ group; preferably L is an alkyl group containing from 4 to 10 carbon atoms, preferably T represents a $-NH-$ group, preferably V represents a $-NH-$ group and m is preferably 1.

In a preferred embodiment, the invention relates to novel affinity ligands of General Formula (XIV):



wherein $A_1, A_2, B_1, B_2, D_1, D_2, p$ and R_1 have the meanings specified above.

In another preferred embodiment, the invention relates to novel affinity ligands of General Formula (XV):



wherein $A_1, A_2, B_1, B_2, D_1, D_2, p$ and R_1 have the meanings specified above and

j is an integer between 2 and 20.

In another preferred embodiment, the invention relates to affinity ligands of General Formula (X), (XI), (XII), (XIII), (XIV) and (XV) wherein D1 and D2 both independently represent a guanidino group, an imidazole group, a primary amino group, a diethylamino group, a trimethylammonium group or a triethylammonium group.

In another preferred embodiment, the invention relates to affinity ligands of General Formula (IX), (X), (XI), (XII), (XIII), (XIV) and (XV) wherein p is 1.



In another preferred embodiment, the invention relates to affinity ligands of General Formula (IX), (X), (XI), (XII), (XIII), (XIV) and (XV) wherein B1 and B2 both independently represent a $-\text{CHCOOH}-(\text{CH}_2)_3-$ group, a $-\text{CHCOOH}-(\text{CH}_2)_4-$ group, a butyl group, a pentyl group or a phenyl group.





In another preferred embodiment, the invention relates to affinity ligands of General Formula (IX), (X), (XI), (XII), (XIII), (XIV) and (XV) wherein A1 and A2 both independently represent a -NH- group.





15 In another preferred embodiment, the invention relates to affinity ligands of General Formula (IX), (X), (XI), (XII) and (XIII) wherein X represents a nitrogen atom.


In another preferred embodiment, the invention relates to affinity ligands of General Formula (XII), (XIII) and (XV) wherein j is between 4 and 10.

20 Preferred affinity ligands according to the invention are:











A valuable group of affinity ligand-matrix conjugates is represented by the General Formula (XXVII):




wherein B_1 , B_2 , D_1 , D_2 , p , M , R_1 , R_2 and R_3 have the meanings hereinbefore specified and j is an integer between 4 and 10.

- 15 An especially valuable group of affinity ligand support matrices is represented by the General Formula (XXVIII):





wherein B₁, B₂, D₁, D₂, p, X, M, R₁, R₂, R₃ and M have the meanings hereinbefore specified.





- 25 Typically, reaction of compounds of General Formula (XXIX):








with 3-propoxy-(1,2-epoxy) derivatised matrices at temperatures between 10°C and 30°C in the presence of an acid binding agent produces novel affinity-ligand matrix conjugates which are of outstanding value in the binding of endotoxin from water, aqueous solutions, proteins, drugs, blood and plasma.

Preferred affinity ligand matrix conjugates according to the invention are









wherein M is as defined above.

10 The invention further covers the use of all such affinity ligand-support matrices in the separation, isolation, purification, quantification, identification and characterisation of endotoxin or analogues, fragments, derivatives thereof and precursors.

Endotoxins are a family of lipopolysaccharides, often abbreviated as LPS, which share a common structure. Endotoxins exist in a number of forms, for example the most significant endotoxin types comprising lipid A attached to a core polysaccharide component which may also be linked to a O-specific chain polysaccharide. The core polysaccharide component may consist of an inner core, an inner core attached to an outer oligosaccharide or an inner core attached to an outer core. Endotoxin is known to be extremely heterogeneous, particularly with respect to the O-Specific Chain polysaccharide and the outer core polysaccharide. Since endotoxin is known to be heterogeneous, the term "endotoxin" as used herein includes all naturally occurring forms which comprise lipid A covalently linked to a polysaccharide, including analogues, derivatives, fragments and precursors thereof and all such forms, irrespective of their source, are subject to the claims of this invention.

Furthermore, the invention relates to a method of attaching the novel affinity ligands of General Formulae (IX) as defined above, (X) as defined above, (XII) as defined above, (XV) as defined above and (XXIX) as defined above to carbohydrate or organic polymer matrices by reacting the carbohydrate or organic polymer matrix with an activating agent followed by reaction of the activated matrix with the novel affinity ligand, optionally in the presence of an

acid binding agent. The invention also relates to a method of attaching the novel affinity ligands of General Formulae (XIII) as defined above to carbohydrate or organic polymer matrices by condensation with the matrix. The invention furthermore relates to a method of attaching the novel affinity ligands of General Formulae (IX) as defined above, (X) as defined above, (XII) as defined above, (XV) as defined above and (XXIX) as defined above to metal oxide, glass or silica matrices, optionally coated with an organic polymer by reacting the optionally coated metal oxide, glass or silica matrix with an activating agent followed by reaction of the activated matrix with the novel affinity ligand, optionally in the presence of an acid binding agent. Another embodiment of the invention relates to a method of attaching the novel affinity ligands of General Formulae (XIII) as defined above to metal oxide, glass or silica matrices optionally coated with an organic polymer by condensation with the matrix. In another embodiment the invention relates to a method of attaching novel affinity ligands of General Formula (XI) as defined above, (XIV) as defined above and (XVI - XXVI) as defined above to a matrix of General Formula (VII) as defined above by reacting the novel affinity ligands with the matrix at temperatures between -0°C and 100°C, optionally in the presence of an acid-binding agent. The invention also relates to all the affinity ligand-matrix conjugates, prepared as described in the above methods.

In another embodiment, the invention relates to the use of the affinity ligands according to the invention and the affinity ligand-matrix conjugates according to the invention for the separation, isolation, purification, characterisation, identification or quantification of endotoxin. In another embodiment the invention relates to any process whereby endotoxin containing solutions or liquids are applied to affinity ligand-matrix conjugates according to the invention at a pH in the range 1.0 to 13.0. The invention also relates to a process for the isolation of endotoxin from various fluids such as water, aqueous solutions, body fluids, blood, plasma, solutions of pharmaceutical products, proteins and other compounds of biological origin by carrying out affinity chromatography using as the biospecific ligand a ligand of General Formula (I) as defined above.

Another embodiment of the invention relates to the use of affinity ligands according to the invention and affinity ligand-matrix conjugates comprising such ligands according to the invention for the extracorporeal removal of endotoxin from whole blood or plasma which is taken from a donor and re-infused back into the same donor or another recipient following treatment.

The invention will now be described in further detail with reference to the following Examples. The examples are provided for illustrative purposes and are not to be construed as limiting the scope of the invention in any way.

Example 1

This Example illustrates the synthesis of a typical affinity ligand of General Formula (XIV) defined by the reaction of a halogenoheterocyclic compound of General Formula (IV) with a compound of General Formula (V) and (VI).

A solution of 1 part cyanuric chloride in 10 parts acetone was added dropwise to a stirred solution comprising 2 parts L-arginine in 100 parts water. The mixture was stirred for 2 hours at 0-5°C whereupon the solution was warmed to 30°C and mixing continued for a further 16 hours. The pH was maintained within the range 5.0 - 7.0 throughout by titration with 1M sodium hydroxide solution. The reaction product was precipitated by the addition of solid sodium chloride to a final concentration of 20% (w/v), filtered and dried in-vacuo. TLC analysis (THF/propan-2-ol/water (1:2:1 by vol.) solvent) revealed the presence of a single reaction product (R_f 0.42). The molecular mass of the isolated compound was determined by mass spectroscopy and found to be consistent with a compound comprising cyanuric chloride derivatised with 2 molecules of arginine (calculated M_r = 459.5; molecular ion (+ve FAB) = 460.3, (-ve FAB) = 458.4). The $^1\text{H-NMR}$ spectrum was consistent with a compound containing arginine.

Example 2

This Example illustrates the synthesis of an optionally derivatised support matrix of General Formula (VII).

A solution of 1 part 1,6-diaminohexane in 12 parts water was added to a stirred suspension comprising 29 parts epoxy-activated agarose beads (30 μ mol epoxide groups per g agarose gel) in 48 parts water and stirred for 24 h at 30°C. The amino-hexyl agarose gel was filtered and washed consecutively with 12 x 29 parts water and allowed to drain under gravity on completion of the final wash. Analysis of the resulting amino-hexyl agarose for the presence of primary amines (TNBS assay) and epoxide groups (thiosulphate/sodium hydroxide titration) revealed complete reaction of the epoxide groups with 1,6-diaminohexane.

10 **Example 3**

Example 2 was repeated by replacing 1,6-diaminohexane with 1,4-diaminobutane, 1,5-diaminopentane, 1,7-diaminoheptane, 1,8-diaminooctane, 1,9-diaminononane and 1,10-diaminodecane. In all cases amino-alkyl derivatised agarose matrices were obtained.

15 **Example 4**

This Example illustrates the reaction of an optionally derivatised support matrix of General Formula (VII) with a halogenoheterocyclic compound of general formula (IV).

A solution of 1 part cyanuric chloride in 10 parts acetone was added to a pre-cooled (0-4°C) suspension of 40 parts amino-hexyl agarose matrix prepared according to Example 2 in 40 parts of 0.5M potassium phosphate buffer, pH 7.0. The mixture was stirred for 1 hour at 0-4°C, filtered and washed consecutively with 5 x 40 parts of a solution comprising 1 part acetone and 1 part water, 5 x 40 parts water, 5 x 40 parts of a solution comprising 1 part acetone and 1 part water and 10 x 40 parts water. Analysis of the resulting dichlorotriazine-activated agarose matrix for the presence of amines (TNBS assay) and release of chloride ions following treatment with 1M sodium hydroxide revealed complete reaction of cyanuric chloride with the primary amino groups on the aminohexyl agarose matrix.

30 **Example 5**

Example 4 was repeated by replacing the product prepared according to Example 2 with the products prepared according to Example 3. Analysis of the

resulting dichlorotriazine-activated agarose matrices for the presence of amines (TNBS assay) and release of chloride ions following treatment with 1M sodium hydroxide revealed in all cases the complete reaction of cyanuric chloride with the primary amino groups on the aminoalkyl agarose matrix.

5

Example 6

This Example illustrates the reaction of the product prepared according to Example 4 with a compound of General Formula (VI) and a compound of General Formula (V) to produce affinity ligand-matrix conjugates of General Formula (III). All solutions were prepared with pyrogen free water.

10

One part arginine was added to a suspension containing 35 parts of the product prepared according to Example 4 in 105 parts of 0.1 M sodium carbonate buffer, pH 10.25. The mixture was stirred for 24 hours at 30°C, filtered and washed consecutively with 12 x 35 parts of 0.1 M sodium carbonate buffer, pH 10.25 and allowed to drain under gravity. This material was re-slurried in 105

15

parts of 0.1 M sodium carbonate buffer, pH 10.25.

20

One part N- ϵ -t-BOC-L-lysine was added to the agarose slurry and the mixture agitated for 72 hours at 85°C. The suspension was filtered and washed consecutively with 12 x 35 parts of water and allowed to drain under gravity. The mixture was resuspended in 35 parts 0.1M trifluoroacetic acid, stirred for 1 hour at 20°C, filtered and washed consecutively with 3 x 35 parts methanol, 12 x 35 parts of water and allowed to drain under gravity. This procedure is required to remove the t-BOC protecting group.

Examples 7 to 14

Table 1 gives further examples of the synthesis of novel affinity ligand-matrix conjugates of the invention which were prepared by the above method but with arginine replaced by the amine compound listed in Column II of Table 1, and N- ϵ -t-BOC-L-lysine replaced by the amine compound listed in Column III of Table 1. The number of the Example is given in Column I of Table 1.

30

Table 1

I	II	III
7	Arginine	Arginine
8	Arginine	Ammonia
9	N-im-Trityl-L-histidine	Arginine
10	N-ε-t-BOC-L-lysine	N-im-Trityl-L-histidine
11	N-ε-t-BOC-L-lysine	N-ε-t-BOC-L-lysine
12	N-im-Trityl-L-histidine	N-im-Trityl-L-histidine
13	N-im-Trityl-L-histidine	Ammonia
14	N-ε-t-BOC-L-lysine	Ammonia

Example 15

This Example illustrates the ability of affinity ligand-matrix conjugates of General Formula (III) to bind endotoxin from water.

Affinity ligand-matrix conjugate (150 µl) prepared according to Example 6 was added to water (1.5 ml) containing *Escherichia coli* #055:B5 endotoxin (1.5×10^4 EU) and agitated for 1 hour at 20°C. The sample was centrifuged and the supernatant assayed for the presence of endotoxin by the Limulus Amoebocyte Lysate Chromogenic Test. Only 0.1 EU/ml (equivalent to 10 pg/ml) was detected in the supernatant indicating greater than 99.99% removal of endotoxin from water.

Examples 16 to 23

Table 2 gives further examples of the ability of novel affinity ligand-matrix conjugates of the invention to bind endotoxin. The procedure described above was performed except that the affinity ligand-matrix conjugate was synthesised according to the Example number given in Column II of Table 2, the amount of endotoxin remaining in the supernatant (EU/ml) is given in Column III of Table 2 and the amount of endotoxin adsorbed (%) by the affinity ligand-matrix conjugate is given in Column IV of Table 2. The number of the Example is given in Column I of Table 2.

Table 2

I	II	III	IV
16	7	1.2	99.99
17	8	1.1	99.99
18	9	1.1	99.99
19	10	7.3	99.93
20	11	0.4	>99.99
21	12	0.1	>99.99
22	13	1.4	99.99
23	14	1.0	99.99

Example 24

This Example illustrates the ability of affinity ligand-matrix conjugates of General Formula (III) to isolate endotoxin from protein containing solutions contaminated with endotoxin.

Affinity ligand-matrix conjugate (150 μ l) prepared according to Example 6 was added to water (1.5 ml) containing *Escherichia coli* #055:B5 endotoxin (1.5×10^4 EU) and human serum albumin (15 mg) and agitated for 1 hour at 20°C. The sample was centrifuged and the supernatant assayed for the presence of endotoxin by the Limulus Amoebocyte Lysate Chromogenic Test which had been calibrated to detect endotoxin in the presence of 10 mg/ml human serum albumin. Endotoxin at a concentration of 50 EU/ml was detected in the supernatant indicating 99.5% removal of endotoxin from a solution containing 10 mg/ml human serum albumin.

Examples 25 to 32

Table 3 gives further examples of the ability of novel affinity ligand-matrix conjugates of the invention to isolate endotoxin from protein containing solutions contaminated with endotoxin. The procedure described above was performed except that the affinity ligand-matrix conjugate was synthesised according to the Example number given in Column II of Table 3, the amount of endotoxin remaining in the supernatant (EU/ml) is given in Column III of Table 3 and the amount of endotoxin adsorbed (%) by the affinity ligand-matrix conjugate is given

in Column IV of Table 3. The number of the Example is given in Column I of Table 3.

Table 3

	I	II	III	IV
5	25	7	790	92.1
	26	8	169	98.3
	27	9	127	98.7
	28	10	81	99.2
	29	11	90	99.1
10	30	12	277	97.2
	31	13	54	99.5
	32	14	66	99.3

Example 33

15 This Example illustrates the capacity of novel affinity ligand-matrix conjugates of General Formula (III) to bind endotoxin in the presence of protein. Affinity ligand-matrix conjugate (150 μ l) prepared according to Example 6 was added to water (1.5 ml) containing *Escherichia coli* #055:B5 endotoxin (7.5×10^3 EU to 1.5×10^5 EU) and human serum albumin (15 mg) and agitated for 1 hour
 20 at 20°C. The samples were centrifuged and the supernatants assayed for the presence of endotoxin by the Limulus Amoebocyte Lysate Chromogenic Test which had been calibrated to detect endotoxin in the presence of 10 mg/ml human serum albumin. The total amount of endotoxin present in the uptake mixture and the amount of endotoxin adsorbed is given in Table 4.

Table 4

	Total Endotoxin (EU)	Endotoxin Bound (%)
25	7.5×10^3	>99.9
	1.5×10^4	99.5
	3×10^4	99.7
30	4.5×10^4	99.6
	7.5×10^4	99.6
	1.5×10^5	99.3

These results demonstrate 1 g of novel affinity ligand-matrix conjugate of
 35 General Formula (III) is able to bind 1×10^6 EU (100 μ g endotoxin) in the


presence of 10mg/ml human serum albumin with an extraction efficiency of greater than 99%.

Example 34

This Example illustrates the ability of novel affinity ligand-matrix
5 conjugates of General Formula (III) to bind endotoxin in the presence of protein
and buffer of varying ionic strength. Affinity ligand-matrix conjugate (150 μ l)
prepared according to Example 6 was added to water (1.5 ml) containing
Escherichia coli #055:B5 endotoxin (7.5×10^4 EU), human serum albumin (15
mg) and PBS buffer (0 to 200 mM) and agitated for 1 hour at 20°C. The samples
10 were centrifuged and the supernatants assayed for the presence of endotoxin
by the Limulus Amoebocyte Lysate Chromogenic Test which had been calibrated
to detect endotoxin in the presence of 10 mg/ml human serum albumin. More
than 99% of the endotoxin present was adsorbed for all concentrations of PBS
buffer investigated. These results demonstrate that novel affinity ligand-matrix
15 conjugates of General Formula (III) are able to bind endotoxin with high
efficiency and independently of ionic strength or the presence of protein.

CLAIMS

1. An affinity ligand-matrix conjugate comprising the matrix and, conjugated thereto by the group Z, a ligand having the general formula (I):



5 wherein one X is N and the other X is N, CCl or CCN;

A₁ and A₂ are each independently O, S or N-R₁ and R₁ is H, C₁₋₆ alkyl, C₁₋₆ hydroxyalkyl, benzyl or β-phenylethyl;

B₁ and B₂ are each independently an optionally substituted hydrocarbon linkage containing from 1 to 10 carbon atoms;

10 D₁ is H or a primary amino, secondary amino, tertiary amino, quaternary ammonium, imidazole, guanidino or amidino group; and

D₂ is a secondary amino, tertiary amino, quaternary ammonium, imidazole, guanidino or amidino group; or

B₂-D₂ is -CHCOOH-(CH₂)₃₋₄-NH₂; and

15 p is 0 or 1.

2. A conjugate according to claim 1, wherein A₁ and A₂ are each independently N-R₁, wherein R₁ is as defined in claim 1.

3. A conjugate according to claim 2, wherein A₁ and A₂ are each NH.

4. A conjugate according to any preceding claim, wherein B₁ and B₂ are each
20 independently -CHCOOH-(CH₂)₁₋₄- or a divalent ethyl, propyl, 2-hydroxypropyl, butyl,
pentyl, hexyl, phenyl, naphthyl or cyclohexyl group.

5. A conjugate according to claim 4, wherein B₁ and B₂ are each independently -CHCOOH-(CH₂)₃₋₄- or a divalent butyl, pentyl or phenyl group.

6. A conjugate according to any preceding claim, wherein D_1 is H, amino, imidazolyl, guanidino, aminidino, trimethylammonium, triethylammonium, dimethylamino, diethylamino, methylamino or ethylamino.

7. A conjugate according to any preceding claim, wherein D_2 is imidazolyl, 5 guanidino, aminidino, trimethylammonium, triethylammonium, dimethylamino, diethylamino, methylamino or ethylamino.

8. A conjugate according to any preceding claim, wherein p is 1.

9. A conjugate according to any preceding claim, wherein each X is N.

10. A conjugate according to any preceding claim, wherein Z is

10



wherein T is O, S or NR_2 and R_2 is H or C_{1-6} alkyl;

V is O, S, $-COO-$, $CONH$, $NHCO$, $-PO_3H-$, NH -arylene- $SO_2-CH_2-CH_2-$ or $N-R_3$

15 and R_3 is H or C_{1-6} alkyl;

L is an optionally substituted hydrocarbon linkage containing from 2 to 20 carbon atoms; and

m is 0 or 1.

11. A conjugate according to claim 10, wherein R_2 and R_3 are each H.

20 12. A conjugate according to claim 10 or claim 11, wherein T is O or NH.

13. A conjugate according to claim 12, wherein T is NH.

14. A conjugate according to any of claims 9 to 13, wherein m is 1 and L is a divalent butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl or dodecyl group.

15. A conjugate according to any of claims 10 to 14, wherein m is 1 and V is O, 25 $-CONH-$, $-NHCO-$ or $N-R_3$.

16. A conjugate according to claim 15, wherein V is O or NH.

17. A conjugate according to claim 16, wherein V is NH.


18. A conjugate according to claim 16 or claim 17, wherein $Z-M$ (M being the matrix) is $-NH-(CH_2)_{4-10}-NH-M$.

30 19. A conjugate according to claim 18, selected from any of formulae XXX to XXXX.

20. A conjugate according to any preceding claim, wherein the matrix is optionally tresyl-activated, sulphonyl chloride-activated, tosyl-activated, vinyl sulphone-activated or epoxy-activated.

21. An affinity ligand of formula (1) as defined in any of claims 1 to 9, wherein Z is 35 a functional group capable of reaction with a solid matrix.

22. A ligand according to claim 21, wherein the matrix is as defined in claim 20.
23. A ligand according to claim 22, wherein Z is F, Cl, Br or I.
24. A ligand according to claim 22, wherein Z is -NH-(CH₂)₂₋₂₀-NH₂.
25. A ligand according to claim 22, wherein Z is -NH-(CH₂)₂₋₂₀-(CO)₀₋₁-OH.
- 5 26. A ligand according to claim 22, wherein Z is



wherein T, L, V and m are as defined in any of claims 10 to 17;

- 10 26. R₁₀ is CO, CH₂, NH-CH₂- or -S-CH₂-; and
R₁₁ is H or C₁₋₆ alkyl.
27. A ligand according to claim 23, wherein Z is Cl, and each X is N.
28. A ligand according to claim 24, wherein each X is N.
29. A ligand according to claim 22, selected from any of formulae XVI to XXVI.
- 15 30. The use of an affinity ligand or conjugate according to any preceding claim, for the separation, isolation, purification, characterisation, identification or quantification of an endotoxin.
31. Use according to claim 30, for the removal of an endotoxin from water or an aqueous solution, body fluid, blood, plasma, solution of pharmaceutical products,
- 20 32. Use according to claim 31, which comprises using the conjugate for the extracorporeal removal of endotoxin from whole blood or plasma, prior to re-infusion into the donor or another recipient.
33. Use according to any of claim 30 to 32, wherein the endotoxin originates from
- 25 34. Use according to claims 30 to 35, wherein an endotoxin-containing solution or liquid is applied to the conjugate at a pH of 1.0 to 13.0.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 00/01759

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 B01J20/32 C02F1/28 C07K1/22 C07D251/50 C07D251/54
C07D403/12 C07D251/70 A61M1/36 G01N33/50

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 B01J C02F C07K G01N B01D C07D A61M

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, EPO-Internal, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 10887 A (NOVO NORDISK) 27 March 1997 (1997-03-27) page 5, line 21 -page 6, line 31; claims 1,2,24,28,30,32,34	1-4, 6-18, 20-28
X	BURTON N P ET AL: "DESIGN OF NOVEL AFFINITY ADSORBENTS FOR THE PURIFICATION OF TRYPSIN-LIKE PROTEASES" JOURNAL OF MOLECULAR RECOGNITION,GB,HEYDEN & SON LTD., LONDON, vol. 5, 1992, pages 55-68, XP000654088 ISSN: 0952-3499 page 58, paragraph 3 - paragraph 4 page 60; table 1 page 66; table 5 page 65; figure 4	1-3, 6-18, 20-24, 27,28

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

11 September 2000

Date of mailing of the international search report

18/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hilgenga, K

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/01759

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 41185 A (PERSEPTIVE BIOSYSTEMS) 19 December 1996 (1996-12-19) claims 1,14,34,36 -----	1,6,7, 30-33
A	GB 2 092 470 A (TANABE SEIYAKU) 18 August 1982 (1982-08-18) page 2, line 27 - line 32 page 13, line 15 - line 30 page 1, line 48 page 1, line 13 - line 19 -----	1-4, 7-18, 20-22, 24,25, 30,31,33
A	EP 0 350 004 A (TANABE SEIYAKU) 10 January 1990 (1990-01-10) page 3, line 26 - line 56 page 4, line 43 - line 47; claim 1 -----	1,30,31, 33,34
A	US 4 229 537 A (L.T. HODGINS) 21 October 1980 (1980-10-21) column 5, line 1 - line 10 -----	
A	US 4 491 660 A (R.L. GENDRICH) 1 January 1985 (1985-01-01) claim 1 -----	
A	EP 0 431 593 A (DAICEL CHEMICAL INDUSTRIES) 12 June 1991 (1991-06-12) -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/01759

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9710887	A	27-03-1997	AU 702901 B AU 6924296 A CA 2232626 A CN 1200680 A EP 0959975 A JP 11512442 T	11-03-1999 09-04-1997 27-03-1997 02-12-1998 01-12-1999 26-10-1999
WO 9641185	A	19-12-1996	NONE	
GB 2092470	A	18-08-1982	DE 3204544 A FR 2499429 A JP 1588983 C JP 2014325 B JP 57183712 A JP 1656109 C JP 3007681 B JP 63118301 A SE 461505 B SE 8200715 A US 4381239 A	02-09-1982 13-08-1982 19-11-1990 06-04-1990 12-11-1982 13-04-1992 04-02-1991 23-05-1988 26-02-1990 11-08-1982 26-04-1983
EP 350004	A	10-01-1990	DE 68915214 D DE 68915214 T DK 169448 B JP 2193072 A JP 2524406 B US 5342785 A	16-06-1994 22-09-1994 31-10-1994 30-07-1990 14-08-1996 30-08-1994
US 4229537	A	21-10-1980	CA 1124190 A DE 2965887 D EP 0012751 A IL 56604 A JP 62059124 B JP 55500309 T WO 7900609 A	25-05-1982 25-08-1983 09-07-1980 31-05-1982 09-12-1987 29-05-1980 23-08-1979
US 4491660	A	01-01-1985	NONE	
EP 431593	A	12-06-1991	AT 107537 T AT 115888 T CA 2022924 A CA 2031599 A CN 1049296 A, B CN 1053046 A, B DE 69010124 D DE 69010124 T DE 69015334 D DE 69015334 T DK 418517 T DK 431593 T EP 0418517 A ES 2058701 T ES 2068975 T GR 3015505 T JP 3238004 A KR 9301609 B US 5032281 A US 5136032 A	15-07-1994 15-01-1995 10-02-1991 08-06-1991 20-02-1991 17-07-1991 28-07-1994 13-10-1994 02-02-1995 17-08-1995 19-09-1994 01-05-1995 27-03-1991 01-11-1994 01-05-1995 30-06-1995 23-10-1991 06-03-1993 16-07-1991 04-08-1992